arXiv:hep-ph/0605093v1 9 May 2006

QCD condensates with flavor SU(3) symmetry breaking

Seung-il Nam* and Hyun-Chul Kim†

Department of Physics and Nuclear Physics & Radiation Technology Institute (NuRI),
Pusan National University, Busan 609-735, Republic of Korea

(Dated: May 2006)

We investigate the effects of flavor SU(3)-symmetry breaking on the QCD condensates, based on the nonlocal effective chiral action from the instanton vacuum, focusing on the quark-gluon mixed one. We employ two different parameterizations for the dependence of the momentum-dependent dynamical quark mass on the strange current quark mass. We obtain the ratios of the mixed condensates: $[\langle\bar{s}\sigma_{\mu\nu}G^{\mu\nu}s\rangle/\langle\bar{u}\sigma_{\mu\nu}G^{\mu\nu}u\rangle]^{1/5}=0.87$. It turns out that the dimensional parameter $m_0^2=\langle\bar{q}\sigma_{\mu\nu}G^{\mu\nu}q\rangle/\langle\bar{q}q\rangle=1.60\sim1.92\,\mathrm{GeV}^2$.

PACS numbers: 11.15.Tk,14.40.Aq
Keywords: Condensates, Instanton vacuum, Flavor SU(3)-symmetry breaking

I. INTRODUCTION

In this proceeding [1], we discuss the effect of flavor SU(3)-symmetry breaking focusing on the quark-gluon mixed condensate, based on the instanton liquid model for the QCD vacuum [1, 2]. The model was later extended by introducing the current quark masses [3, 4]. Since we are interested in the effect of explicit flavor SU(3)-symmetry breaking, we follow the formalism in Ref. [3]. Though the mixed quark-gluon condensate was already studied in the instanton vacuum [5], explicit SU(3)-symmetry breaking was not considered. Hence, we extend the work of Ref. [5], focussing on the effect of flavor SU(3)-symmetry breaking on the QCD vacuum. Furthermore, we take into account two different parameterizations for the dependence of the dynamical quark mass on the current quark mass m_f so that we can examine the effects of the current quark mass very in detail. We observed that, with a proper choice of the m_f dependence of the dynamical quark mass [6], the gluon condensate is independent of m_f [7]. The corresponding results are summarized as follows: The ratio $[\langle\bar{s}\sigma_{\mu\nu}G^{\mu\nu}s\rangle/\langle\bar{u}\sigma_{\mu\nu}G^{\mu\nu}u\rangle]^{1/5}=0.87$, $m^2_{0,\mathrm{u}}=1.60\,\mathrm{GeV}^2$, and $m^2_{0,\mathrm{s}}=1.84\,\mathrm{GeV}^2$, with isospin symmetry assumed. This proceeding is based on our previous work of Ref. [7] in which the gluon and quark condensates are were also discussed in addition to the mixed one.

II. FORMALISM

Now, we consider the mixed condensate, $\langle\bar{q}\sigma_{\mu\nu}G^{\mu\nu}q\rangle$ in our framework. Actually, the local operator inside this condensate corresponds to the quark-gluon interaction of a Yukawa type. However, in the present work, the gluon field strength ($G_{\mu\nu}$) can be expressed in terms of the quark-instanton interaction [5]. First, the one flavor quark and one instanton interaction can be rewritten as a function of space-time coordinates x and color orientation matrix U:

$$Y_{\pm,1}(x,U)=\int\frac{d^4k}{(2\pi)^4}\frac{d^4p}{(2\pi)^4}[2\pi\rho F(k\rho)][2\pi\rho F(p\rho)]e^{-ix\cdot(k-p)}\left[U^{\alpha}_{i'}(U^{j'}_{\beta})^{\dagger}\epsilon^{ii'}\epsilon_{jj'}\right]\left[i\psi^{\dagger}(k)_{\alpha i}\frac{1\pm\gamma_5}{2}\psi(p)^{\beta j}\right],\tag{1}$$

where the form factor, $F(k\rho)$, is defined as follows:

$$F(k\rho)=2t\left[I_0(t)K_1(t)-I_1(t)K_0(t)-\frac{1}{t}I_1(t)K_1(t)\right].\tag{2}$$

$\bar{\rho}$ is set to be $1/600\ \mathrm{MeV}^{-1}$. We assume the δ-function-type instanton distribution. We define then the field strength $G^a_{\mu\nu}$ in terms of the instanton configuration as a function of $I(\bar{I})$ position and orientation matrix U:

$$G^a_{\pm\mu\nu}(x,x',U)=\frac{1}{2}\left[\lambda^aU\lambda^bU^{\dagger}\right]G^b_{\pm\mu\nu}(x'-x).\tag{3}$$

---

*Electronic address: sinam@pusan.ac.kr
†Electronic address: hchkim@pusan.ac.kr

[1] This proceeding is prepared for the international workshop, "Hadrons at finite density" (HFD06) at YITP in Japan 20∼22 Feb 2006.

$G^b_{\pm\mu\nu}(x'-x)$ stands for the field strength consisting of a certain instanton configuration. Using Eqs. (1) and (3), we define the field strength in terms of the quark-instanton interaction:

$$\hat{G}^a_{\pm\mu\nu} = \frac{iN_c M}{4\pi\bar{\rho}^2}\int d^4x \int dU G^a_{\pm\mu\nu}(x,x',U)Y_{\pm,1}(x,U) \tag{4}$$

Following the method in Ref. [5], we finally obtain the quark-gluon mixed condensate as follows:

$$\langle\bar{q}\sigma_{\mu\nu}G^{\mu\nu}q\rangle = 2N_c\bar{\rho}^2\int\frac{d^4k_1}{(2\pi)^4}\int\frac{d^4k_2}{(2\pi)^4}\frac{\sqrt{M(k_1)M(k_2)}G(k_1,k_2)N(k_1,k_2)}{[k_1^2+[m_f+M(k_1)]^2][k_2^2+[m_f+M(k_2)]^2]}, \tag{5}$$

where $G(k_1,k_2)$ and $N(k_1,k_2)$ are defined as follows:

$$\begin{aligned} G(k_1,k_2) &= 32\pi^2\left[\frac{K_0(t)}{2}+\left\{\frac{4K_0(t)}{t^2}+\left(\frac{2}{t}+\frac{8}{t^3}\right)K_1(t)-\frac{8}{t^4}\right\}\right], \\ N(k_1,k_2) &= \frac{1}{(k_1-k_2)^2}\left[8k_1^2k_2^2-6(k_1^2+k_2^2)k_1\cdot k_2+4(k_1\cdot k_2)^2\right] \end{aligned} \tag{6}$$

with $t=|k_1-k_2|\bar{\rho}$. If we consider for arbitrary N_f the mixed condensate, the situation may be somewhat different from the cases of the gluon and quark condensates. We note that, though the mixed condensate has been calculated for the case of $N_f=1$, the same formula of Eq. (5) still holds for each flavor with arbitrary N_f as discussed previously [5].

As indicated in Refs. [3, 8], the dynamical quark mass is also a function of the current quark mass. Thus, we redefine the dynamical quark mass as a function of momentum and current quark mass, m_f:

$$M(k)\rightarrow M(k,m_f)=M_0 f(m_f)F^2(k\bar{\rho}), \tag{7}$$

where we set M_0 to be 350 MeV. In order to take into account the dependence of the dynamical quark mass on m_f, we introduce the m_f-dependent correction factor $f(m_f)$ [3, 8]. We consider here two different parameterizations for $f(m_f)$ (A and B) of $f(m_f)$ to see theoretical ambiguities:

$$f_A(m_f)=\sqrt{1+\frac{m_f^2}{c_A^2}}-\frac{m_f}{c_A},\quad f_B(m_f)=1+\frac{c_B}{M_0},\quad f_C(m_f)=1. \tag{8}$$

The correction factor $f_A(m_f)$ is derived by Pobylitsa [6] by expanding the quark propagator in the instanton background in terms of the large N_c limit. The parameter c_A is set to 0.198 GeV. The m_f correction factor $f_B(m_f)$, was introduced in Ref. [3] by using the saddle-point equation and its expansion in the current quark mass m_f. We use the value of $c_B=-0.5\,m_f$ as proposed in Ref. [3]. f_C is for the M without m_f dependence.

III. NUMERICAL RESULTS

In the right panel of Fig. 1, we draw the results of the quark-gluon mixed condensate as functions of the m_f. The curves of the mixed condensate decrease as m_f increases. We also list the values of the mixed condensate for the up and strange quarks in Table. I.

	A	B	C
$\langle\bar{u}\sigma_{\mu\nu}G^{\mu\nu}u\rangle$	-481^5	-483^5	-484^5
$\langle\bar{s}\sigma_{\mu\nu}G^{\mu\nu}s\rangle$	-418^5	-475^5	-483^5

TABLE I: Quark-gluon mixed condensates for $m_u=5$ and $m_s=150$ MeV [MeV^5].

Now, we consider the effect of flavor SU(3)-symmetry breaking by calculating the ratios between the nonstrange condensates and the strange ones. As already discussed, the ratio of the gluon condensates remains unity for the m_f correction factor f_A. The calculated ratios are given by; $[\langle\bar{s}\sigma_{\mu\nu}G^{\mu\nu}s\rangle/\langle\bar{u}\sigma_{\mu\nu}G^{\mu\nu}u\rangle]^{1/5}=0.87\sim1.00$. Being compared to Refs. [13, 14, 15, 16, 17], our results are in good agreement each other.

We take into account a dimensional quantity m_0^2 defined as the ratio between the mixed and quark condensates:

$$m_0^2=\langle\bar{q}\sigma_{\mu\nu}G^{\mu\nu}q\rangle/\langle\bar{q}q\rangle, \tag{9}$$

which is an important input for general QCD sum rule calculations. We draw m_0^2 in the left panel of Fig. 1 as a function of m_f and list the values of m_0^2 in Table. II in which the quark condensates are computed in the same framework.

The value of m_0^2 increases as m_f does, which implies that the mixed condensate is less sensitive to the m_f than the quark condensate. The values of m_0^2 are in the range of $1.84\,\mathrm{GeV}^2$ for the strange quark and of $1.60\,\mathrm{GeV}^2$ for the up quark. Thus, the strange $m_{0,s}^2$ turns out to be larger than the nonstrange $m_{0,u}^2$ by about 15 %.



FIG. 1: Left: Quark-gluon mixed condensate as a function of the current quark mass m_f. Right: m_0^2 as a function of the current quark mass. The solid curve draws the mixed condensate for $f_A(m_f)$ in Eq. (8), the dashed one corresponds to that without any $f(m_f)$, and the dotted one plots that for $f_B(m_f)$. We used $c_A = 0.198$ GeV and $c_B = -0.5\,m_f$ GeV.

	A	B	C
$[\langle\bar{s}\sigma_{\mu\nu}G^{\mu\nu}s\rangle/\langle\bar{u}\sigma_{\mu\nu}G^{\mu\nu}u\rangle]^{1/5}$	0.87	0.98	1.00
$m_{0,u}^2 = \langle\bar{u}\sigma_{\mu\nu}G^{\mu\nu}u\rangle/\langle\bar{u}u\rangle$	1.60 GeV^2	1.60 GeV^2	1.60 GeV^2
$m_{0,s}^2 = \langle\bar{s}\sigma_{\mu\nu}G^{\mu\nu}s\rangle/\langle\bar{s}s\rangle$	1.84 GeV^2	1.89 GeV^2	1.92 GeV^2

TABLE II: The ratios of the condensates for the different types of the m_f correction factors. $m_u = 5$ MeV and $m_s = 150$ MeV are used.

IV. SUMMARY AND CONCLUSION

In the present work, we investigated the various QCD vacuum condensates within the framework of the instanton liquid model, emphasizing the effects of flavor SU(3)-symmetry breaking. The modified improved action elaborated by Musakhanov was used for this purpose. In the modified improved action, the current quark mass appeared explicitly in the denominator of the quark propagator as well as in the dynamical quark mass. Thus, we were able to take into account the current quark mass effects to the QCD condensates. In order to consider the m_s dependence of the dynamical quark mass, we employed two different types of the correction factors, f_A and f_B (f_C for the case without the correction). f_A arises from the resummation of the QCD planar loops in the large N_c limit [6], while f_B is a simple parameterization of the current quark mass correction suggested by Musakhanov [3].

The mixed condensates were calculated with these correction factors. The results were consistent with those from other model calculations as well as phenomenological values. In particular, The ratios of the condensates between the strange and up quarks were also investigated: $[\langle\bar{s}\sigma_{\mu\nu}G^{\mu\nu}s\rangle/\langle\bar{u}\sigma_{\mu\nu}G^{\mu\nu}u\rangle]^{1/5}$ and $\langle\bar{q}\sigma_{\mu\nu}G^{\mu\nu}q\rangle/\langle\bar{q}q\rangle$. It turned out that the results are again compatible with other theoretical calculations. The dimensional quantity m_0^2 was also studied: $m_{0,u}^2 = 1.6\,\mathrm{GeV}^2$ and $m_{0,s}^2 = 1.84\,\mathrm{GeV}^2$. In general, the quark and mixed condensates decrease as the current quark mass increases. However, the m_0^2 increases as the current quark mass does, which indicates that the mixed condensate is less sensitive to the current quark mass than the quark condensate. More details can be found in Ref. [7].

Acknowledgments

The present work is supported by the Korean Research Foundation (KRF–2003–070–C00015). The authors are grateful to M. M. Musakhanov and Y. Kwon for fruitful discussions. S. I. N. would like to thank A. Hosaka and

RCNP for the support during the YITP workshop "Hadrons at finite density" at Kyoto, Japan 20∼22 Feb 2006.

---

[1] D. Diakonov and V. Y. Petrov,Nucl. Phys. B **272**, 457 (1986).
[2] D. Diakonov, Prog. Part. Nucl. Phys. **51**, 173 (2003).
[3] M. Musakhanov, Eur. Phys. J. C **9**, 235 (1999).
[4] M. Musakhanov, Nucl. Phys. A **699**, 340 (2002).
[5] M . V. Polyakov and C. Weiss, Phys. Lett. B **387**, 841 (1996).
[6] P. V. Pobylitsa, Phys. Lett. B **226**, 387 (1989).
[7] S. i. Nam and H. Ch-. Kim, arXiv:hep-ph/0605041.
[8] H.-Ch. Kim, M. M. Musakhanov and M. Siddikov, Phys. Lett. B **633**, 201 (2006).
[9] T. Schafer and E. V. Shuryak, Rev. Mod. Phys. **70**, 323 (1998).
[10] M. Jamin, Phys. Lett. B **538**, 71 (2002).
[11] H. G. Dosch, M. Jamin and S. Narison, Phys. Lett. B **220**, 251 (1989).
[12] A. Di Giacomo and Y. A. Simonov, Phys. Lett. B **595**, 368 (2004).
[13] M. Beneke and H. G. Dosch, Phys. Lett. B **284**, 116 (1992).
[14] K. Aladashvili and M. Margvelashvili, Phys. Lett. B **372**, 299 (1996).
[15] V. M. Khatsimovsky, I. B. Khriplovich and A. R. Zhitnitsky, Z. Phys. C **36**, 455 (1987).
[16] A. A. Ovchinnikov and A. A. Pivovarov, Sov. J. Nucl. Phys. **48**, 721 (1988) [Yad. Fiz. **48**, 1135 (1988)].
[17] V. M. Braun and A. Lenz, Phys. Rev. D **70**, 074020 (2004).